

Drug Testing

Essential Elements
Helen Harberts MA JD
Chief Probation Officer (ret)
Chief Deputy DA (ret)

Drug and Alcohol Testing

Adult Best Practices Standards Volume 2, Section 7

Testing must be:

- Frequent
- Random
- Long enough
- Wide breadth
- Witnessed collection
- Valid specimens
- Accurate and reliable testing
- Rapid results
- Participant contracts
- Due process

Various sample types exist

- Hair
- Oral fluids
- Breath (ETOH)
- Electronic transdermal
- Urine-most common

Each sample has limits, strengths and weaknesses. You must know them all.

Frequent testing

- Urine testing: at least twice per week (7 day week) throughout program. EtG & EtS too
- Monitor via transdermal devices or oral fluids in a more comprehensive manner. Requires more frequency due to shortened detection window.
- The more frequent the testing the better the outcomes (within limits). *Certainty of detection matters!*

Random

- Random and unpredictable.
- Nights and weekends and holidays are the same probability as M-F.
- Rapid production of sample after notification (reduces adulteration and dilutes)
- You need to mix things up. If you are being tested a minimum of two times per week, you should have a 2 in 7 chance of being tested each day (28%). Consider a 30 day cycle.

Duration

- Both drug and alcohol testing is uninterrupted throughout the drug court tenure.
- It is a form of continued assessment and supports recovery.
- It is critical to measure treatment progress and monitor for lapses.
- While other contacts and services are reduced over time, testing is NOT.

Breadth of testing

- Broad panels of tests are required to monitor changes in drug use patterns. Testing for only arrest drug, or perceived drugs of choice may produce unearned negative tests.
- Monitor both drugs and alcohol constantly.
- Be aware of limits on opiate tests, in particular
- There are new drugs being used all the time, watch for them.

Witnessed Collection

- Direct observation of urine testing is required.
- The person supervising the test must be trained to detect tampering, or adulteration of tests.
- Absent very unusual circumstances, alternative testing by others is not allowed.
- If there is a suspicion of urine testing fraud, use a secondary test such as oral fluid as well.

Valid Specimens

- Specimens must be examined for dilution (creatinine testing) and adulteration.
- Temperature strips if no direct observation
- Adulteration strips for instant tests
- No Band-Aids, washed hands, etc.

WHY must we directly witness urine production?

Quick Fix

- Quick Fix Whizzinator Pack
- One Whizzinator
- One Quick Fix
- \$149.95



<http://www.quickfixurine.com/>

Squat and Cough: You don't do it...
this is what gets past you.





Oral Fluids?

- [Saliva Swab Drug Test](#)
- Mouthwash

- [Oral Clear Gum \(Saliva\)**](#)
Neutralizing Gum \$90
99.9% Success Rate!



Unsupervised Drug Test

		
Instant Clean Add-Itive* 8 ml. \$75 \$50 99.9% Success Rate!	Spike Additive* 2 (TWO) 1.5 ML Vials \$125 99.9% Success Rate!	Sub-Solution* (UniSex) Synthetic Urine \$75\$55 100% Success Rate!

Addiction makes folks do strange things....this, of course, is an example of an attempted urine substitution.



Substitution



NOT ACTUAL SIZE !

The WEDGE Kit Includes :

1. The WEDGE
2. Fill Bottle
3. Temperature Strip
4. Tube Clamp
5. Clean Swab
6. Clean Gloves
7. Closed Cell Foam

And Easy To Read Directions.

THE URINATOR Substitution



“The urinator the ultimate urine testing device only \$149.95. A digitally temperature controlled unit that is reusable, reliable and by far the most superior product on the market.”


PRE-FILLED KIT

New!



Lil' Whizz Kit

New, from the makers of the legendary Whizzinator! A Non-refillable, use and dispose 3oz belt!
Everything you need right out of the box. Works every time. Clean, Safe, Dependable, and Toxin Free Synthetic Urine.



This is the original design of CUPID and uses the patented Flow Restrictors to actuate the flow urine. The only difference between CUPID and Original SheWhizz is the size of the bottle. CUPID was designed around a 2 oz bottle. And is exactly double the size of SheWhizz's 1 fl oz bottle. Made for those times when your golden shower fantasy needs to be TWICE the fun! NOW WITH NEW TEXTURED SURFACE!

Full Kit contains: 1- CUPID Device, 1- 10ml Syringe, 1- 3-pack of Flow Restrictors, 1- 3 fl oz bottle of Laboratory Grade Synthetic Urine

\$ 89.99

CUPID now comes pre-filled with dehydrated, synthetic urine and fully incased in latex. You simply clip off the sealed hose caps, fill with warm water and shake.

https://youtu.be/ZJQLMM_G-YY







Accurate and Reliable Testing Procedures (chemistry)

- Valid, reliable and legally admissible testing procedures
- CHAIN OF CUSTODY with paper trail for evidence.
- Confirmation upon denial of a positive screen
- Chromatography confirmations
- Industry standard cut off levels are followed to determine positive tests and to interpret testing for new use or changes in use patterns.

Accurate and Reliable Testing Procedures (chemistry)

- Learn and apply distinctive rules for different substances
 - Cannabis
 - Clean out time, no passive inhalation, etc.
 - Oxy
 - Poor bonding on many opiate instant tests
 - EtG & EtS
 - Cut off levels, and the Nyquil excuses

Rapid Results

- Test results back within 48 hours of sample collection, including confirmation.
- Timing, like certainty, impacts behavior modification. You need results, *fast*.

Participant contracts

- Clear contract of expectations for drug testing
 - Dilutes, missed tests, insufficient tests, adulterated, or failure to provide a sample will be addressed with a sanction.
 - All tests will be directly observed by a staff person.
 - Rights to challenge, and a hearing, or a retest, at their expense if they are wrong.
 - Responsibilities: call in, respond quickly,
 - Value of telling truth first.
 - Cannabis rules "up front", synthetic use, etc.
 - Violations of the "no association" rule. Etc.
